

EPIMORPHISMS OF 3-MANIFOLD GROUPS

MICHEL BOILEAU AND STEFAN FRIEDL

ABSTRACT. Let $f: M \rightarrow N$ be a proper map between two aspherical compact orientable 3-manifolds with empty or toroidal boundary. We assume that N is not a closed graph-manifold. Suppose that f induces an epimorphism on fundamental groups. We show that f is homotopic to a homeomorphism if one of the following holds: either for any finite-index subgroup Γ of $\pi_1(N)$ the ranks of Γ and of $f_*^{-1}(\Gamma)$ agree, or for any finite cover \tilde{N} of N the Heegaard genus of \tilde{N} and the Heegaard genus of the pull-back cover \tilde{M} agree.

1. INTRODUCTION

Let $f: M \rightarrow N$ be a map between two 3-manifolds. (Here and throughout the paper all 3-manifolds are understood to be connected, compact and orientable.) We say f is *proper* if $f(\partial M) \subset \partial N$. We say f is a π_1 -*epimorphism* if the induced map $f_*: \pi_1(M) \rightarrow \pi_1(N)$ is an epimorphism. Finally f is called a *degree-one map* if it is proper and if the induced map on homology $f_*: H_3(M, \partial M; \mathbb{Z}) \rightarrow H_3(N, \partial N; \mathbb{Z})$ is an isomorphism. It is well-known, see e.g. [He76, Lemma 15.12], that a degree-one map is a π_1 -epimorphism.

Given a map $f: M \rightarrow N$ between two 3-manifolds that is either a π_1 -epimorphism or a degree-one map it is a long-standing question to find out what extra conditions ensure that f is in fact homotopic to a homeomorphism. For example, a celebrated theorem of M. Gromov and W. Thurston says that if $f: M \rightarrow N$ is a degree-one map between two hyperbolic 3-manifolds of the same volume, then f is homotopic to a homeomorphism, [Th79, Chapter 6], see also [BCG95]. This result has been generalized in many directions, see e.g. [Som95, Der10, Der12], by using the Gromov simplicial volume [Gr82].

For an aspherical 3-manifold M the volume $\text{Vol}(M)$ is defined as the sum of the volumes of the hyperbolic pieces of the geometric decomposition of M . In the following we say that a map $f: M \rightarrow N$ between manifolds is a \mathbb{Z} -*homology equivalence*, if f induces isomorphisms of all homology groups. We recall the following theorem of B. Perron-P. Shalen and P. Derbez.

Theorem 1.1. *Let $f: M \rightarrow N$ be a map between two closed, aspherical 3-manifolds with the same volume. If for any finite covering $\tilde{N} \rightarrow N$ (not necessarily regular)*

2010 *Mathematics Subject Classification.* 57M10, 57M27.

Key words and phrases. 3-manifolds, degree-one maps, Heegaard genus.

the induced map $\tilde{f}: \tilde{M} \rightarrow \tilde{N}$ is a \mathbb{Z} -homology equivalence, then f is homotopic to a homeomorphism.

Remark.

- (1) The case of closed graph manifolds was dealt with by B. Perron-P. Shalen [PS99], see also [AF11].
- (2) P. Derbez [Der03] stated the theorem for Haken 3-manifolds that are not graph manifolds. The proof of P. Derbez applies to all closed 3-manifolds, that are not graph manifolds, for which the Geometrization Conjecture is known. In particular by the work of G. Perelman the theorem applies to all aspherical 3-manifolds that are not graph manifolds.
- (3) The original theorem of P. Derbez is formulated in terms of the simplicial volume, but the simplicial volume for 3-manifolds agrees, up to a fixed constant, with the volume. See e.g. [BP92] for details.
- (4) The conclusion of the theorem does not hold without the assumption on the simplicial volume for in [BW96] an example of two closed aspherical 3-manifolds M and N is given, such that for any finite covering $\tilde{N} \rightarrow N$ the induced map $\tilde{f}: \tilde{M} \rightarrow \tilde{N}$ is a \mathbb{Z} -homology equivalence, but such that $\text{Vol}(M) > \text{Vol}(N)$.

In the following recall that a subgroup $\Gamma \subset \pi$ is called *subnormal* if there exists a chain of subgroups $\pi = \Gamma_0 \supset \Gamma_1 \supset \dots \supset \Gamma_k = \Gamma$, such that each Γ_i is normal in Γ_{i-1} . Given a finitely generated group π we denote by $\text{rk}(\pi)$ its rank, i.e. the smallest cardinality of a generating set of π . If $f: M \rightarrow N$ is a π_1 -epimorphism, then given any finite-index subgroup Γ of $\pi_1(N)$ the map f_* induces an epimorphism $f_*^{-1}(\Gamma) \rightarrow \Gamma$, in particular the inequality

$$\text{rk}(f_*^{-1}(\Gamma)) \geq \text{rk}(\Gamma)$$

holds. Our first theorem says that equality holds for all finite-index Γ 's only if f is homotopic to a homeomorphism.

Theorem 1.2. *Let $f: M \rightarrow N$ be a proper map between two aspherical 3-manifolds with empty or toroidal boundary. We assume that N is not a closed graph manifold. If f is a π_1 -epimorphism and if for every finite-index subnormal subgroup Γ of $\pi_1(N)$ the equality*

$$\text{rk}((f_*)^{-1}(\Gamma)) = \text{rk}(\Gamma)$$

holds, then f is homotopic to a homeomorphism.

One of the key ingredients of the proof of Theorem 1.2 is the virtual fibering theorem, stated as Theorem 2.4. Only 3-manifolds with empty or toroidal boundary can be (virtually) fibered. This explains the restriction to the case of 3-manifolds with empty or toroidal boundary.

Before we can state our second theorem we need to introduce a few more definitions.

- (1) We say that a covering $p: \widehat{N} \rightarrow N$ of a manifold N is *subregular* if the covering p can be written as a composition of coverings $p_i: N_i \rightarrow N_{i-1}$, $i = 1, \dots, k$ with $N_k = \widehat{N}$ and $N_0 = N$, such that each p_i is regular.
- (2) Given a 3-manifold M we denote by $h(M)$ the minimal number of one-handles in a handle-composition of M with one zero-handle. If M is closed, then $h(M)$ equals the *Heegaard genus*, i.e. the minimal genus of a Heegaard surface of M . If M has non-empty boundary, $h(M)$ is smaller or equal to the minimal genus of a Heegaard splitting of the 3-manifold triad $(M; \emptyset, \partial M)$ as defined by A. Casson and C. Mc Gordon [CG87], see also M. Scharlemann [Scha02].

Our second theorem gives another variation on Theorem 1.1.

Theorem 1.3. *Let $f: M \rightarrow N$ be a proper map between two aspherical 3-manifolds with empty or toroidal boundary. We assume that N is not a closed graph manifold. If f is a π_1 -epimorphism and if for every finite subregular cover \widetilde{N} of N and induced cover \widetilde{M} the inequality $h(\widetilde{M}) \leq h(\widetilde{N})$ holds, then f is homotopic to a homeomorphism.*

Using the examples given in [Li013] and [SW07] of closed aspherical 3-manifolds with arbitrarily large discrepancy between the Heegaard genus and the rank of the fundamental group, it is possible to build examples of π_1 -epimorphism $f: M \rightarrow N$ between two aspherical closed 3-manifolds where the inequality $h(M) < h(N)$ can be arbitrarily large (see Section 4). However to the best of our knowledge it is still not known whether for every degree-one map $f: M \rightarrow N$ between two aspherical 3-manifolds the inequality $h(M) \geq h(N)$ holds.

The following result, which is basically a consequence of the proof of Theorem 1.3, shows that the inequality always holds virtually, even for a proper π_1 -epimorphism.

Proposition 1.4. *Let $f: M \rightarrow N$ be a proper map between two aspherical 3-manifolds with empty or toroidal boundary. We assume that N is not a closed graph manifold. If f is a π_1 -epimorphism, then there exists a finite subregular cover \widetilde{N} of N such that the induced cover \widetilde{M} satisfies the inequality $h(\widetilde{M}) \geq h(\widetilde{N})$. See Conjecture 4.2.*

Remark. In Theorems 1.2, 1.3 and in Proposition 1.4 we excluded the case that N is a closed graph manifold. What we really need for the statements to hold is that N is a 3-manifold that is virtually fibered. By the Virtual Fibration Theorem of I. Agol [Ag08, Ag13] P. Przytycki–D. Wise [PW14, PW12] and D. Wise [Wi09, Wi12a, Wi12b] any aspherical 3-manifold that is *not* a closed graph manifold is virtually fibered. The three results also apply if N is a closed graph manifold that is virtually fibered, e.g. if N is a Sol-manifold or if N carries a Riemannian metric of nonpositive sectional curvature [Liu013]. We do not know though whether our results hold for aspherical graph manifolds that are not virtually fibered. We refer to [LW93, Ne96] for more examples of graph manifolds that are not virtually fibered.

This paper is organized as follows. In Section 2 we provide the proof of Theorem 1.2. The proof of Theorem 1.3 has many formal similarities with the proof of Theorem 1.3.

In Section 3 we will point out how to modify the proof of Theorem 1.2 to obtain a proof of Theorem 1.3.

Acknowledgments. Work on this paper was supported by the SFB 1085 ‘Higher Invariants’ at the Universität Regensburg funded by the Deutsche Forschungsgemeinschaft (DFG). We also wish to thank the referee for helpful feedback.

2. THE RANK OF FUNDAMENTAL GROUPS AND π_1 -EPIMORPHISMS

2.1. Proof of Theorem 1.2 in the fibered case. Let M be a 3-manifold. Throughout this paper we identify $H^1(M; \mathbb{Z})$ with $\text{Hom}(\pi_1(M), \mathbb{Z})$. We say a primitive class $\phi \in H^1(M; \mathbb{Z}) = \text{Hom}(\pi_1(M), \mathbb{Z})$ is *fibered* if there exists a fibration $p: M \rightarrow S^1$ such that the induced map $p_*: \pi_1(M) \rightarrow \pi_1(S^1) = \mathbb{Z}$ coincides with ϕ . It follows from Stallings’ theorem [St62] (together with the resolution of the Poincaré conjecture) that ϕ is a fibered class if and only if $\ker(\phi: \pi_1(M) \rightarrow \mathbb{Z})$ is finitely generated.

Proposition 2.1. *Suppose that $f: M \rightarrow N$ is a proper π_1 -epimorphism between two 3-manifolds. We assume that there exists a primitive class $\phi \in H^1(N; \mathbb{Z})$ such that $\phi \in H^1(N; \mathbb{Z})$ and $f^*\phi \in H^1(M; \mathbb{Z})$ are fibered. If $\text{rk}(\ker(f^*\phi)) = \text{rk}(\ker(\phi))$, then $f_*: \pi_1(M) \rightarrow \pi_1(N)$ is an isomorphism.*

Proof. We denote by F the fiber of ϕ and we denote by E the fiber of $f^*\phi$. These are compact orientable surfaces.

We consider the following commutative diagram of short exact sequences

$$\begin{array}{ccccccc} 1 & \longrightarrow & \pi_1(E) & \longrightarrow & \pi_1(M) & \xrightarrow{\phi \circ f_*} & \mathbb{Z} \longrightarrow 0 \\ & & \downarrow f_* & & \downarrow f_* & & \downarrow = \\ 1 & \longrightarrow & \pi_1(F) & \longrightarrow & \pi_1(N) & \xrightarrow{\phi} & \mathbb{Z} \longrightarrow 0. \end{array}$$

Since $f_*: \pi_1(M) \rightarrow \pi_1(N)$ is an epimorphism it follows that the map on the left is also an epimorphism. By our hypothesis we have

$$\text{rk}(\pi_1(E)) = \text{rk}(\ker(f^*\phi)) = \text{rk}(\ker(\phi)) = \text{rk}(\pi_1(F)).$$

We make the following claim.

Claim. The induced map $\pi_1(E) \rightarrow \pi_1(F)$ is an isomorphism.

First consider the case that M is closed. In that case E is also closed. Together with the observation that $f_*: \pi_1(E) \rightarrow \pi_1(F)$ is an epimorphism and from the observation that $\text{rk}(\pi_1(E)) = \text{rk}(\pi_1(F))$ we deduce that F is also closed and that the map $f_*: \pi_1(E) \rightarrow \pi_1(F)$ is an isomorphism.

Now we consider the case that M has boundary. Then also E has boundary. Since f is a proper map the manifold N , and thus also F have boundary. Thus $f_*: \pi_1(E) \rightarrow \pi_1(F)$ is an epimorphism between free groups of the same rank, thus it is already an isomorphism. This concludes the proof of the claim.

An elementary argument using the above commutative diagram shows that the induced map $f_*: \pi_1(M) \rightarrow \pi_1(N)$ is an isomorphism. \square

2.2. Ranks and fibers.

Proposition 2.2. *Let M be a 3-manifold. We write $\pi = \pi_1(M)$. Furthermore let $\phi \in H^1(M; \mathbb{Z}) = \text{Hom}(\pi, \mathbb{Z})$ be a fibered class. We write*

$$\pi_n = \ker(\pi_1(M) \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}/n).$$

Let p be a prime. For any n we have

$$\dim_{\mathbb{F}_p}(H_1(\pi_n; \mathbb{F}_p)) \leq \text{rk}(\pi_n) \leq 1 + \text{rk}(\ker(\phi)).$$

Furthermore, there exists an n such that

$$\dim_{\mathbb{F}_p}(H_1(\pi_n; \mathbb{F}_p)) = \text{rk}(\pi_n) = 1 + \text{rk}(\ker(\phi)).$$

Proof. We denote by F the fiber of the fibration corresponding to ϕ and we write $\Gamma = \pi_1(F)$. We can identify π with a semidirect product $\langle t \rangle \rtimes_{\mu} \Gamma$ in such a way that $\phi(t) = 1$ and $\phi|_{\Gamma}$ is trivial. Here $\mu: \Gamma \rightarrow \Gamma$ denotes the corresponding automorphism of Γ , which is just given by the monodromy action on $\Gamma = \pi_1(F)$.

Given any n we have

$$\pi_n = \langle t^n \rangle \rtimes \Gamma.$$

It follows that

$$\dim_{\mathbb{F}_p}(H_1(\pi_n; \mathbb{F}_p)) \leq \text{rk}(\pi_n) \leq 1 + \text{rk}(\Gamma) = 1 + \text{rk}(\ker(\phi)).$$

Now let p be a prime. The homology group $H_1(\Gamma; \mathbb{F}_p)$ is finite. Thus there exists an n such that μ_*^n acts like the identity on $H_1(\Gamma; \mathbb{F}_p)$. It follows that

$$\begin{aligned} \text{rk}(\pi_n) &= \text{rk}(\langle t^n \rangle \rtimes_{\mu} \Gamma) = \text{rk}(\langle t \rangle \rtimes_{\mu^n} \Gamma) \geq H_1(\langle t \rangle \rtimes_{\mu^n} \Gamma; \mathbb{F}_p) \\ &= \text{rk}(\mathbb{Z} \oplus H_1(\Gamma; \mathbb{F}_p)) \\ &= 1 + \text{rk}(\Gamma) = 1 + \text{rk}(\ker(\phi)). \end{aligned}$$

Here we used that for the fundamental group of the orientable compact surface F we have $\text{rk}(\pi_1(F)) = \dim_{\mathbb{F}_p}(H_1(F; \mathbb{F}_p))$. \square

2.3. The rank gradient and fiberedness. Let π be a group and $\phi: \pi \rightarrow \mathbb{Z}$ be a homomorphism. We write

$$\pi_n = \ker(\pi \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}/n).$$

Following [La05] and [DFV14] we refer to

$$\text{rg}(\pi, \phi) := \liminf_{n \rightarrow \infty} \frac{1}{n} \text{rk}(\pi_n)$$

as the *rank gradient* of (π, ϕ) .

Theorem 2.3. *Let M be a 3-manifold and let $\phi: \pi_1(M) \rightarrow \mathbb{Z}$ be a non-trivial homomorphism. Then the following two statements are equivalent:*

- (1) $\phi \in \text{Hom}(\pi_1(M), \mathbb{Z}) = H^1(M; \mathbb{Z})$ is fibered,
- (2) $\text{rg}(\pi_1(M), \phi) = 0$.

Here the implication (1) \Rightarrow (2) is an immediate consequence of Proposition 2.2. The implication (2) \Rightarrow (1) is the main result of [DFV14] (see also [De16]). The proof in [DFV14] builds on the fact that twisted Alexander polynomials detect fibered manifolds [FV12], that proof in turn relies on the recent results of D. Wise [Wi09, Wi12a, Wi12b].

2.4. Proof of Theorem 1.2. Before we provide a proof of Theorem 1.2 we recall the Virtual Fibered Theorem of Agol [Ag08, Ag13], Przytycki–Wise [PW14, PW12] and Wise [Wi09, Wi12a, Wi12b]. (We refer to [AFW15] for the precise references for the Virtual Fibered Theorem.)

Theorem 2.4. *Any prime 3-manifold with empty or toroidal boundary that is not a closed graph manifold admits a finite cover that is fibered.*

Now we are finally in a position to prove Theorem 1.2.

Proof of Theorem 1.2. Suppose that $f: M \rightarrow N$ is a proper π_1 -epimorphism between two aspherical 3-manifolds with empty or toroidal boundary. We assume N is not a closed graph manifold. We suppose that the following condition holds:

- (*) Given any finite-index subnormal subgroup Γ of $\pi_1(M)$ the equality

$$\text{rk}(f_*^{-1}(\Gamma)) = \text{rk}(\Gamma)$$

holds.

We will show that f is homotopic to a homeomorphism. Since N is a prime 3-manifold with empty or toroidal boundary that is not a closed graph manifold it follows from Theorem 2.4 that N admits a finite-index regular covering $p: \tilde{N} \rightarrow N$ such that \tilde{N} admits a primitive fibered class $\phi \in H^1(\tilde{N}; \mathbb{Z}) = \text{Hom}(\pi_1(\tilde{N}), \mathbb{Z})$.

We denote by \tilde{M} the finite-cover of M corresponding to $\pi_1(\tilde{M}) = f_*^{-1}(\pi_1(\tilde{N}))$. Given $n \in \mathbb{N}$ we denote by \tilde{M}_n the cover of \tilde{M} corresponding to the epimorphism

$$\pi_1(\tilde{M}) \xrightarrow{\phi \circ f_*} \mathbb{Z} \rightarrow \mathbb{Z}/n.$$

Similarly we denote by \tilde{N}_n the cover of \tilde{N} corresponding to the epimorphism

$$\pi_1(\tilde{N}) \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}/n.$$

Evidently $\pi_1(\tilde{N}_n)$ is a subnormal subgroup of $\pi_1(\tilde{N})$ and we have $\pi_1(\tilde{M}_n) = f_*^{-1}(\pi_1(\tilde{N}_n))$. By our assumption (*) we have

$$\text{rk}(\pi_1(\tilde{M}_n)) = \text{rk}(\pi_1(\tilde{N}_n)) \text{ for all } n.$$

Since ϕ is fibered it follows from Theorem 2.3 that $\text{rg}(\pi_1(\tilde{N}), \phi) = 0$. By the above observation we have $\text{rg}(\pi_1(\tilde{M}), f_*\phi) = 0$. It is a consequence of Theorem 2.3 that $f_*\phi$ is a fibered class of \tilde{M} .

Similarly, it is a consequence of the above observations and of Proposition 2.2 that $\text{rk}(\ker(f^*\phi)) = \max \{\text{rk}(\pi_1(\widetilde{M})_n) - 1 \mid n \in \mathbb{N}\} = \max \{\text{rk}(\pi_1(\widetilde{N})_n) - 1 \mid n \in \mathbb{N}\} = \text{rk}(\ker(\phi))$.

It is now a consequence of Proposition 2.1 that $f_*: \pi_1(\widetilde{M}) \rightarrow \pi_1(\widetilde{N})$ is an isomorphism.

Next we show that $f_*: \pi_1(M) \rightarrow \pi_1(N)$ is already an isomorphism. We consider the following diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \pi_1(\widetilde{M}) & \longrightarrow & \pi_1(M) & \longrightarrow & \pi_1(M)/\pi_1(\widetilde{M}) \longrightarrow 1 \\ & & \cong \downarrow f_* & & \downarrow f_* & & \cong \downarrow f_* \\ 1 & \longrightarrow & \pi_1(\widetilde{N}) & \longrightarrow & \pi_1(N) & \longrightarrow & \pi_1(N)/\pi_1(\widetilde{N}) \longrightarrow 1. \end{array}$$

This diagram consists of maps between pointed sets (i.e. sets with distinguished elements). The top and bottom sequence are exact in the category of pointed sets. The left and the rightmost maps are isomorphisms of pointed sets. The diagram evidently commutes. Thus it follows from the five-lemma that the middle map is also an isomorphism.

Summarizing we showed that $f_*: \pi_1(M) \rightarrow \pi_1(N)$ is an isomorphism. Since $f: (M, \partial M) \rightarrow (N, \partial N)$ is a proper map, it follows from work of Waldhausen [Wa68, Corollary 6.5][He76, Theorem 13.16] and from Mostow-Prasad rigidity [Mo68, Pr73] that f is homotopic to a homeomorphism. \square

3. THE HEEGAARD GENUS AND π_1 -EPIMORPHISMS

3.1. Proof of Theorem 1.3. In this section we will provide the proof of Theorem 1.3. We start out with the following well-known lemma.

Lemma 3.1. *Let M be a manifold. Then the following hold:*

- (1) $\text{rk}(\pi_1(M)) \leq h(M)$.
- (2) *If M is a fibered 3-manifold with connected fiber F , then*

$$h(M) \leq b_1(F) + 1.$$

Proof. By definition we can endow M with a handle decomposition with one zero-handle and $h(M)$ one-handles. It is thus clear that the free group on $h(M)$ generators surjects onto $\pi_1(M)$. Therefore $\text{rk}(\pi_1(M)) \leq h(M)$. This concludes the proof of (1).

Now we turn to the proof of (2). If M is fibered with fiber F , then we can write $M = F \times [0, 4]/(x, 0) \sim (\varphi(x), 4)$. First we consider the case that M is closed. Pick two closed disks $P \subset \Sigma$ and $Q \subset \Sigma$ with $P \cap Q = \emptyset$ and $\varphi(Q) \cap P = \emptyset$. It is straightforward to see that

$$\overline{(F \setminus (P \cup Q))} \times \{0\} \cup \partial P \times [0, 2] \cup \overline{(F \setminus (P \cup \varphi(Q)))} \times \{2\} \cup \partial \varphi(Q) \times [2, 4]$$

splits M into two handlebodies of genus $2g(F) + 1 = b_1(F) + 1$. It follows that we can endow M with a handle-decomposition with one zero-handle and $2g + 1 = b_1(F) + 1$ one-handles. Now we consider the case that M has non-empty boundary. We pick a

closed disk $P \subset F$. Then the image X of $F \times [0, 1] \cup P \times [1, 3] \cup F \times [3, 4]$ in M is homeomorphic to the handle body $F \times [-1, 1]$ together with one extra one-handle. Put differently, X be written as a handlebody with one zero-handle and $b_1(F) + 1$ one-handles. We can view F as obtained from the disk by attaching $b_1(F)$ one-handles. It is now straightforward to see that M can be built out of X by attaching $b_1(F)$ two-handles. It follows that $h(M) \leq b_1(F) + 1$. \square

The following proposition plays the rôle of Proposition 2.2.

Proposition 3.2. *Let M be a 3-manifold. Furthermore let $\phi \in H^1(M; \mathbb{Z}) = \text{Hom}(\pi_1(M), \mathbb{Z})$ be a primitive fibered class with corresponding fiber F . Given $n \in \mathbb{N}$ we denote by M_n the cover of M corresponding to $\ker(\pi_1(M) \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}/n)$. For any n we have*

$$h(M_n) \leq b_1(F) + 1.$$

Furthermore, there exists an n such that

$$h(M_n) = b_1(F) + 1.$$

Remark. If M is closed and hyperbolic, then Souto [Sou08, Theorem 1.1] showed that there exists an m such that for every $n \geq m$ we have $h(M_n) = 2g(F) + 1 = b_1(F) + 1$.

Proof. We pick a fibration $p: M \rightarrow S^1$ with fiber F and monodromy $\varphi: F \rightarrow F$ that corresponds to the given primitive fibered class ϕ . Since ϕ is primitive the fiber F is connected. For any n the fibration p gives rise to a fibration $p_n: M_n \rightarrow S^1$ with fiber F and monodromy φ^n . The first statement is now a consequence of Lemma 3.1.

By Proposition 2.2 there exists an n such that

$$\text{rk}(\pi_1(M_n)) = 1 + \text{rk}(\pi_1(F)).$$

It follows from Lemma 3.1 that

$$1 + \text{rk}(\pi_1(F)) = \text{rk}(\pi_1(M_n)) \leq h(M_n) \leq 1 + b_1(F) = 1 + \text{rk}(\pi_1(F)).$$

Thus we obtain that $h(M_n) = b_1(F) + 1$. \square

Let M be a 3-manifold and let $\phi: \pi_1(M) \rightarrow \mathbb{Z}$ be an epimorphism. We write M_n for the finite cyclic cover corresponding to

$$\ker(\pi_1(M) \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}/n).$$

Following [La06] we refer to

$$\text{hg}(M, \phi) := \liminf_{n \rightarrow \infty} \frac{1}{n} (2h(M_n) - 2)$$

as the *Heegaard gradient* of (M, ϕ) . The following proposition plays the rôle of Theorem 2.3.

Proposition 3.3. *Let M be a 3-manifold and let $\phi: \pi_1(M) \rightarrow \mathbb{Z}$ be an epimorphism. Then the following two statements are equivalent:*

- (1) $\phi \in \text{Hom}(\pi_1(M), \mathbb{Z}) = H^1(M; \mathbb{Z})$ is fibered,

$$(2) \text{ hg}(M, \phi) = 0.$$

Here the implication (1) \Rightarrow (2) is an immediate consequence of Proposition 3.2. For the implication (2) \Rightarrow (1) note that if $\text{hg}(M, \phi) = 0$, then it follows from Lemma 3.1 that $\text{rg}(M, \phi) = 0$, which in turn implies by Theorem 2.3 that ϕ is fibered.

Remark. The implication (2) \Rightarrow (1) was first proved by M. Lackenby [La06, Theorem 1.11] for closed hyperbolic 3-manifolds (see also [Ren10, Ren14] for extensions). The proof of M. Lackenby is significantly harder than the proof provided in [DFV14] since the latter proof makes use of the results of D. Wise [Wi09, Wi12a, Wi12b] to simplify many arguments.

The proof of Theorem 1.3 is now almost entirely identical to the proof of Theorem 1.2, we only need to replace the study of ranks of subnormal finite-index subgroups of $\pi_1(M)$ and $\pi_1(N)$ by the Heegaard genera of the corresponding subregular finite-index covers of M and N . In particular we need to replace Proposition 2.2 by Proposition 3.2 and we need to replace Theorem 2.3 by Proposition 3.3. We leave the straightforward and dull details to the reader.

3.2. Proof of Proposition 1.4. Let $f: M \rightarrow N$ be a proper π_1 -epimorphism map between two aspherical 3-manifolds with empty or toroidal boundary. We assume that N is not a closed graph manifold. As in the proof of Theorem 1.2 there exists a finite subregular cover \tilde{N} of N which is fibered with fiber F and which satisfies

$$\text{rk}(\pi_1(\tilde{N})) = 1 + 2 \text{rk}(\pi_1(F)).$$

We denote by \tilde{M} the induced cover of M . The map f gives rise to a proper map $f: \tilde{M} \rightarrow \tilde{N}$, which induces an epimorphism of fundamental groups. It follows from Lemma 3.1 that

$$h(\tilde{M}) \geq \text{rk}(\pi_1(\tilde{M})) \geq \text{rk}(\pi_1(\tilde{N})) = 1 + \text{rk}(\pi_1(F)) = 1 + b_1(F) \geq h(\tilde{N}).$$

This concludes the proof of Proposition 1.4.

4. EXAMPLES

The goal of this short section is to show the existence of π_1 -epimorphisms $f: M \rightarrow N$ between two aspherical closed 3-manifolds where the inequality $h(M) < h(N)$ can be arbitrarily large.

Proposition 4.1. *Given any $n \in \mathbb{N}$ there exist closed orientable aspherical 3-manifolds M_n and N_n such that there is a π_1 -epimorphism $f: M_n \rightarrow N_n$ while $h(N_n) > h(M_n) + n$.*

Proof. Given $n \in \mathbb{N}$, there exists a closed aspherical 3-manifold N_n such that $h(N_n) - \text{rk}(\pi_1(N_n)) \geq n + 3$. The manifold N_n can be hyperbolic or have a non-trivial JSJ with a hyperbolic piece by [Li013], or it can be a graph manifold by [SW07].

Given such a manifold N_n there is a π_1 -epimorphism $f: X_n \rightarrow N_n$ where X_n is the connected sum of $\text{rk}(\pi_1(N_n))$ copies of $S^1 \times S^2$. The manifold X_n can be obtained by taking the double of a handlebody H_n of genus $\text{rk}(\pi_1(N_n))$. Then the Heegaard surface ∂H_n is unique up to isotopy by [Wa68b]. Let $k \subset X_n$ be a knot lying on ∂H_n and which meets essentially the boundary of each meridian disk of H_n . Since k can be put in a 1-bridge position with respect to ∂H_n , by [Do91, Theorem 1.6] the exterior $E(k) = X_n \setminus \mathcal{N}(k)$ is a compact irreducible 3-manifold. By construction $h(E(k)) = \text{rk}(\pi_1(N_n)) + 1$, see [Do91, Remark (2)]. Let $W(k)$ be any Whitehead double of k . Then the exterior $E(W(k))$ is obtained from $E(k)$ by gluing along $\partial E(k)$ the exterior W of the Whitehead link. Since $h(W) = 2$, by amalgamating the Heegaard splittings of $E(k)$ and W one gets a Heegaard splitting for $E(W(k))$ of genus $h(E(k)) + h(W) - 1 = h(E(k)) + 1$, see [La04]. It follows that $h(E(W(k))) \leq h(E(k)) + 1 = \text{rk}(\pi_1(N_n)) + 2$. By [Ha82] there are only finitely many boundary slopes on $\partial E(W(k))$. This implies in particular that for almost all slopes α on $\partial E(W(k))$ the Dehn fillings of $E(W(k))$ produces a closed aspherical 3-manifold M . Moreover there is a degree-one map $g: M \rightarrow N_n$, since $W(k)$ is null-homotopic in X_n , see [BW96]. Therefore there is a π_1 -surjective map $f \circ g: M \rightarrow N_n$ while $h(M) \leq h(E(W(k))) \leq \text{rk}(\pi_1(N_n)) + 2 \leq h(N_n) - (n + 1)$. \square

To the best of our knowledge the following conjecture remains open:

Conjecture 4.2. *If there is a degree-one map $f: M \rightarrow N$ between two aspherical 3-manifolds, does the inequality $h(M) \geq h(N)$ hold?*

REFERENCES

- [Ag08] I. Agol, *Criteria for virtual fibering*, J. Topol. 1 (2008), no. 2, 269–284.
- [Ag13] I. Agol, *The virtual Haken conjecture*, with an appendix by I. Agol, D. Groves and J. Manning, Documenta Math. 18 (2013), 1045–1087.
- [AF11] M. Aschenbrenner and S. Friedl, *Residual properties of graph manifold groups*, Top. Appl. 158 (2011), 1179–1191.
- [AFW15] M. Aschenbrenner, S. Friedl and H. Wilton, *3-manifold groups*, EMS Series of Lectures in Mathematics (2015)
- [BP92] R. Benedetti and C. Petronio, *Lectures on hyperbolic geometry*, Universitext, Springer-Verlag, Berlin (1992)
- [BCG95] G. Besson, G. Courtois and S. Gallot, *Entropies et rigidités des espaces localement symétriques de courbure strictement négative*, Geom. Funct. Anal. 5 (1995), no. 5, 731–799.
- [BW96] M. Boileau and S. Wang, *Non-zero degree maps and surface bundles over S^1* , J. Differential Geom. 43 (1996), no. 4, 789–806.
- [CG87] A. Casson and C. McA. Gordon, *Reducing Heegaard splittings*, Topology and its Applications, 27 (1987), 275–283.
- [De16] J. DeBlois, *Explicit rank bounds for cyclic covers*, Algebr. Geom. Topol. 16 (2016), no. 3, 1343–1371.
- [DFV14] J. DeBlois, S. Friedl and S. Vidussi, *The rank gradient for infinite cyclic covers of 3-manifolds*, Michigan Math. J. 63 (2014), 65–81.
- [Der03] P. Derbez, *A criterion for homeomorphism between closed Haken manifolds*, Algebr. Geom. Topol. 3 (2003), 335–398.

- [Der10] P. Derbez, *Topological rigidity and Gromov simplicial volume*, Comment. Math. Helv. 85 (2010), no. 1, 1–37.
- [Der12] P. Derbez, *Local rigidity of aspherical three-manifolds*, Ann. Inst. Fourier (Grenoble) 62 (2012), no. 1, 393–416.
- [Do91] H. Doll, *A generalized bridge number for links in 3-manifolds*, Math. Ann. 294 (1992), 701–717.
- [FV12] S. Friedl and S. Vidussi, *A Vanishing Theorem for Twisted Alexander Polynomials with Applications to Symplectic 4-manifolds*, J. Eur. Math. Soc. 16 (2013) no. 6, 2027–2041.
- [He76] J. Hempel, *3-Manifolds*, Ann. of Math. Studies, no. 86. Princeton University Press, Princeton, N. J., 1976.
- [Gr82] M. Gromov, *Volume and bounded cohomology*, Publications mathématiques de I.H.S. 56 (1982), 5–99.
- [Ha82] A. Hatcher, *On the boundary slopes of incompressible surfaces*, Pacific J. Math. 99 (1982), 373–377.
- [La04] M. Lackenby, *The Heegaard Genus of Amalgamated 3-Manifolds*, Geom Dedicata 109 (2004), 139–145.
- [La05] M. Lackenby, *Expanders, rank and graphs of groups*, Israel J. Math. 146 (2005), 357–370.
- [La06] M. Lackenby, *Heegaard splittings, the virtually Haken conjecture and Property (τ)*, Invent. math. 164 (2006), 317–359.
- [Li013] T. Li, *Rank and genus of 3-manifolds*, J. Amer. Math. Soc. 26 (2013), 777–829.
- [Liu013] Y. Liu, *Virtual cubulation of nonpositively curved graph manifolds*, J. Topol. 6 (2013), 793–822.
- [LW93] J. Luecke and Y.-Q. Wu, *Relative Euler number and finite covers of graph manifolds*, in: Geometric Topology, pp. 80103, AMS/IP Studies in Advanced Mathematics, vol. 2.1, Amer. Math. Soc., Providence, RI; International Press, Cambridge, MA, 1997.
- [Mo68] G. D. Mostow, *Quasi-conformal mappings in n -space and the rigidity of hyperbolic space forms*, Inst. Hautes Études Sci. Publ. Math. 34 (1968), 53–104.
- [Ne96] W. Neumann, *Commensurability and virtual fibration for graph manifolds*, Topology 39 (1996), 355–378.
- [PS99] B. Perron and P. Shalen, *Homeomorphic graph manifolds: A contribution to the μ constant problem*, Topology Appl. 99 (1999), no. 1, 1–39.
- [Pr73] G. Prasad, *Strong rigidity of \mathbb{Q} -rank 1 lattices*, Invent. Math. 21 (1973), 255–286.
- [PW12] P. Przytycki and D. Wise, *Mixed 3-manifolds are virtually special*, Preprint (2012).
- [PW14] P. Przytycki and D. Wise, *Separability of embedded surfaces in 3-manifolds*, Compositio Mathematica 150 (2014), 1623–1630.
- [Ren10] C. Renard, *Gradients de Heegaard sous-logarithmiques d’une variété hyperbolique de dimension 3 et fibres virtuelles*, Actes du Séminaire Théorie Spectrale et Géométrie de Grenoble 29 (2010-2011), 97–131.
- [Ren14] C. Renard, *Detecting surface bundles in finite covers of hyperbolic closed 3-manifolds*, Trans. Amer. Math. Soc. 366 (2014), no. 2, 979–1027.
- [Scha02] M. Scharlemann, *Heegaard splittings of compact 3-manifolds*, Handbook of geometric topology, ed. by R. Daverman and R. Sherr, 921–953, North-Holland, Amsterdam, 2002.
- [SW07] J. Schultens and R. Weidmann, *On the geometric and the algebraic rank of graph manifolds*, Pacific J. Math. 231 (2007), no. 2, 481–510.
- [Som95] T. Soma, *A rigidity theorem for Haken manifolds*, Math. Proc. Cambridge Philos. Soc. 118 (1995), no. 1, p. 141–160.

- [Sou08] J. Souto, *The rank of the fundamental group of certain hyperbolic 3-manifolds fibering over the circle*, Boileau, Michel (ed.) et al., The Zieschang Gedenkschrift. Geometry and Topology Monographs 14 (2008), 505–518.
- [St62] J. Stallings, *On fibering certain 3-manifolds*, 1962 Topology of 3-manifolds and related topics (Proc. The Univ. of Georgia Institute, 1961) pp. 95–100 Prentice-Hall, Englewood Cliffs, N.J. (1962)
- [Th79] W.P. Thurston, *The geometry and topology of three-manifolds*, Princeton University Mathematics Department (1979).
- [Wa68] F. Waldhausen, *On irreducible 3-manifolds which are sufficiently large*, Ann. of Math. (2) 87 (1968), 56–88.
- [Wa68b] F. Waldhausen, *Heegaard-Zerlegungen der 3-Sphäre*, Topology 7 (1968), 195–203.
- [Wi09] D. Wise, *The structure of groups with a quasi-convex hierarchy*, Electronic Res. Ann. Math. Sci 16 (2009), 44–55.
- [Wi12a] D. Wise, *The structure of groups with a quasi-convex hierarchy*, 189 pages, preprint (2012), downloaded on October 29, 2012 from <http://www.math.mcgill.ca/wise/papers.html>
- [Wi12b] D. Wise, *From riches to RAAGs: 3-manifolds, right-angled Artin groups, and cubical geometry*, CBMS Regional Conference Series in Mathematics,

AIX-MARSEILLE UNIVERSITÉ, CNRS, CENTRALE MARSEILLE, I2M, UMR 7373, 13453 MARSEILLE, FRANCE

E-mail address: `michel.boileau@cmi.univ-mrs.fr`

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, GERMANY

E-mail address: `sfriedl@gmail.com`